

HOW TO RESCUE SOLID CLOSURE

HOLGER BRENNER

1 February 2008

ABSTRACT. We define a closure operation for ideals in a commutative ring which has all the good properties of solid closure (at least in the case of equal characteristic) but such that also every ideal in a regular ring is closed. This gives in particular a kind of tight closure theory in characteristic zero without referring to positive characteristic.

INTRODUCTION

The theory of tight closure, introduced by Hochster and Huneke ([7],[8],[9],[10],[2]), is defined for rings containing a field of positive characteristic making use of the Frobenius endomorphism. The main applications of tight closure are to homological conjectures. For example, with this technique it is quite easy to prove that the invariant ring of an action by a linearly reductive group on a polynomial ring is Cohen-Macaulay. This rests upon three facts of tight closure: colon-capturing, persistence and the fact that every ideal in a regular ring is tightly closed.

There are several ways to extend this notion to rings containing a field of characteristic zero, by declaring that an element belongs to the tight closure if and only if this holds for almost all points with residue class field of positive characteristic, where the relevant data have to be expressed in an algebra over a finitely generated $\mathbb{Z}$ -algebra. Most results from tight closure pass then from the case of positive characteristic to characteristic zero.

The theory of solid closure was introduced by Hochster in [5] in order to give a closure operation defined for every Noetherian ring without referring to positive characteristic. It coincides with tight closure if the ring contains a field of positive characteristic under some mild finiteness conditions, and it contains tight closure also in the case of a field of characteristic zero. However, a computation of Roberts in [12] showed that regular rings of dimension ≥ 3 in characteristic zero are not solidly closed. This “discouraging” result shows that solid closure “does not have the right properties in equal characteristic zero” ([6, Introduction]).

This paper proposes another closure operation, called parasolid closure. Roughly speaking, an element h belongs to the parasolid closure of $(f_1, \dots, f_n) \subseteq R$ (where R is local of dimension d), if the forcing algebra A for these data is (universally) parasolid: this means that every canonical element (given by parameters in R) coming from $H_{\mathfrak{m}}^d(R)$ survive in $H_{\mathfrak{m}}^d(A)$.

We list the main properties of this closure operation (compare [9, Introduction]). The parasolid closure is persistent and it lies inside solid closure. Every ideal in a regular ring is parasolidly closed.

If the ring contains a field, then the contraction from a finite extension belongs to the closure. This is not clear in mixed characteristic, and I have to admit

that our closure operation has no effect on the homological conjectures in mixed characteristic.

Over a field of positive characteristic it coincides with solid closure and hence also with tight closure under some mild conditions. Though the definition of parasolid closure does not refer to positive characteristic, reduction mod p is anyway a useful method to prove results for parasolid closure in equal characteristic zero. The theorem of Briançon-Skoda holds for parasolid closure in equal characteristic zero and we prove this with the help of the finiteness theorem of Hochster ([2, Theorem 8.4.1]). Moreover, parasolid closure contains tight closure in characteristic zero, hence also colon capturing holds for it.

For a complete local Gorenstein ring we also define an ideal which coincides with the tight closure test ideal in positive characteristic.

The content of this paper is as follows. Section 1 gives the concept of a parasolid algebra and some characterizations in positive characteristic and in low dimensions. Section 2 gives criteria for an algebra A over a complete local Gorenstein ring R to be parasolid in terms of the module-sections $\mathrm{Hom}_R(A, R)$ and the order ideal in R . We show that a regular complete local ring is in every parasolid algebra a direct summand.

Section 3 introduce the closure operation and establishes its basic properties. In section 4 we show that every ideal in a regular ring is closed. Rings with this property will be called pararegular. Section 5 is concerned with the relation between solid and parasolid closure and with tight closure in positive characteristic.

In section 6 we look at the intersection of the order ideals of all parasolid forcing algebras. We show that for a complete local Gorenstein ring over a field of positive characteristic this intersection is the same as the tight closure test ideal. In section 7 we give characterizations for pararegular rings in the case of a complete Gorenstein ring.

In section 8 we express the datas which say that a paraclass vanishes in terms of a finitely generated $\mathbb{Z}$ -algebra and equations. This gives relations between the parasolid closure in positive characteristic, in zero characteristic and in mixed characteristic. We show that the equations which define over a field a quotient singularity in dimension two yield also in mixed characteristic examples of pararegular, but non regular rings (this is a kind of reduction to the case of a field). We obtain the Theorem of Briançon-Skoda for parasolid closure in equal characteristic zero by reduction to positive characteristic using Hochster's finiteness Theorem. Furthermore we get from a result of Koh [11] a finite type criterion for universally parasolid algebras.

In section 9 we describe results how parasolid algebras and the parasolid closure behave in the fibers of a family, and derive from this as a special case that the tight closure of an ideal in characteristic zero is contained inside the parasolid closure.

1. PARASOLID ALGEBRAS

Let R denote a local Noetherian ring of dimension d with maximal ideal $\mathfrak{m}$. Due to a theorem of Grothendieck we know that the local cohomology $H_{\mathfrak{m}}^d(R) \neq 0$. Elements $x_1, \dots, x_d \in R$ are called parameters if $V(x_1, \dots, x_d) = V(\mathfrak{m})$. A system of parameters defines via Čech cohomology the element $1/x_1 \cdots x_d \in H_{\mathfrak{m}}^d(R)$, and we call such a cohomology class a *canonical element* or a *paraclass*. If R contains a field K , then such a paraclass is $\neq 0$ (The proof for this is different for positive and zero

characteristic, see [2, Remark 9.2.4]). This is not known for mixed characteristic, but at least there exist paraclases $\neq 0$.

The main ingredient of the theory of solid closure is the concept of a solid algebra. For a domain R , an R -algebra A is called *solid* if there exists a non-trivial R -module homomorphism $\varphi : A \rightarrow R$. If R is a local complete Noetherian domain, this is equivalent with $H_{\mathfrak{m}}^d(A) \neq 0$, see [5, Corollary 2.4]. In the following definition we require the stronger condition that all the paraclases coming from $H_{\mathfrak{m}}^d(R)$ shall not vanish in $H_{\mathfrak{m}}^d(A)$.

Definition 1.1. Let R denote a local Noetherian ring of dimension d and let A denote an R -algebra. A is called *parasolid* if the image of every paraclase $0 \neq c \in H_{\mathfrak{m}}^d(R)$ in $H_{\mathfrak{m}}^d(A)$ does not vanish.

Definition 1.2. Let R denote a Noetherian commutative ring and let A be an R -algebra. We say that A is *parasolid* if $A_{\mathfrak{m}}$ is parasolid over $R_{\mathfrak{m}}$ for every maximal ideal $\mathfrak{m} \subset R$.

We say that A is *universally parasolid* (even if R is not Noetherian) if for every local Noetherian ring $R \rightarrow R'$ the algebra $A \otimes_R R'$ is a parasolid R' -algebra.

Remark 1.3. The condition $c \neq 0$ is imposed only for mixed characteristic, for this is not known in general. The paraclase $c = 1/x_1 \cdots x_d \in H_{\mathfrak{m}}^d(R)$ vanishes in $H_{\mathfrak{m}}^d(A)$ (we will often say that it vanishes in A) if and only if $(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ holds in A for some $k \in \mathbb{N}$.

In the definition of universally parasolid it is enough to consider only complete local rings, since completion does not change the d -th local cohomology. In equal characteristic we have only to look at complete normal domains.

Some examples of parasolid algebras are gathered together in the following proposition.

Proposition 1.4. Let R denote a commutative ring and let A denote an R -algebra. In the following situations A is universally parasolid.

- (i) R is a direct summand of A .
- (ii) A is faithfully flat over R .
- (iii) R contains a field and $R \subseteq A$ is a finite extension (or $R \rightarrow A$ is finite and $\text{Spec } A \rightarrow \text{Spec } R$ is surjective).
- (iv) There exists $A \rightarrow A'$ such that A' is a universally parasolid R -algebra.

Proof. We may assume that R is a local Noetherian ring of dimension d and we have to show that A is parasolid.

(i). Let $A = R \oplus V$. Then the cohomological mapping $H_{\mathfrak{m}}^d(R) \rightarrow H_{\mathfrak{m}}^d(A) = H_{\mathfrak{m}}^d(R) \oplus H_{\mathfrak{m}}^d(V)$ is injective.

(ii). In the flat case we know that $H_{\mathfrak{m}}^d(A) = H_{\mathfrak{m}}^d(R) \otimes_R A$. Since A is faithfully flat, it is also pure and again the mapping is injective.

(iii). Let $x_1, \dots, x_d$ be parameters of R and let $(x_1, \dots, x_d)A \subseteq \mathfrak{n}$ be a maximal ideal of A . Then $x_1, \dots, x_d$ are parameters in $A_{\mathfrak{n}}$ and their paraclase does not vanish.

(iv) is clear. □

Proposition 1.5. Let R denote a local Noetherian ring of dimension d and let $\hat{R}$ be its completion. Then an R -algebra A is parasolid if and only if $A' = A \otimes_R \hat{R}$ is parasolid over $\hat{R}$.

Proof. (Here I have to thank Prof. Storch for a hint) We know that $H_m^d(R) = H_m^d(\hat{R})$ and hence the mapping on the cohomology $H_m^d(A) \rightarrow H_m^d(A') = H_m^d(A) \otimes_R R'$ is injective. Hence it is enough to show that the paraclases coming from R and from $\hat{R}$ are the same up to a unit. Let $x_1, \dots, x_d$ be parameters in $\hat{R}$ and let $\mathfrak{a}$ be the ideal they generate. Since $\mathfrak{a}$ is primary, it is the extended ideal of an ideal $\mathfrak{b}$ in R . The number of minimal generators does not change, hence $\mathfrak{b}$ is also a parameter ideal. Therefore $(x_1, \dots, x_d) = (y_1, \dots, y_d)$, where $y_i \in R$.

Then they define up to a unit the same paraclase. This can be proved by replacing inductively x_i by y_i using [1, Exercises 5.1.10-5.1.15]. $\square$

Over a complete local domain a parasolid algebra is solid, and in positive characteristic these notions coincide.

Lemma 1.6. *Let K denote a field of positive characteristic p and let R denote a local Noetherian K -Algebra of dimension d . Let A be an R -algebra. Then the following are equivalent.*

- (i) *A is a parasolid R -algebra.*
- (ii) *There exists a paraclase $c \in H_m^d(R)$ which does not vanish in $H_m^d(A)$.*
- (iii) *$H_m^d(A) \neq 0$.*

Proof. (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii) are clear. (iii) $\Rightarrow$ (i). Suppose there exists a paraclase $1/(x_1 \cdots x_d)$ which does vanish in $H_m^d(A)$, and let $c = a/(x_1^r \cdots x_d^r)$ be an element in $H_m^d(A)$. The vanishing of the paraclase means that there exists an equation in $A_{x_1 \cdots x_d}$,

$$\frac{1}{x_1 \cdots x_d} = \frac{a_1}{(x_2 \cdots x_d)^k} + \cdots + \frac{a_d}{(x_1 \cdots x_{d-1})^k}.$$

Let $q = p^e \geq r$ and apply the e -th Frobenius to this equation. This shows that also the paraclase $1/(x_1 \cdots x_d)^q$ vanishes, hence also $c = 0$. $\square$

Corollary 1.7. *Let K denote a field of positive characteristic p and let R denote a local Noetherian K -Algebra of dimension d . Let A be an R -algebra. Then the following are equivalent.*

- (i) *A is universally parasolid.*
- (ii) *$A \otimes_R \hat{R}/\mathfrak{q}$ is solid for every minimal prime of the completion $\hat{R}$ (his property is called formally solid, see [5, Definition 3.2]).*

If R is a complete local domain, then A is parasolid if and only if it is universally parasolid.

Proof. (i) $\Rightarrow$ (ii) is clear, so assume that (ii) holds. Then from [5, Theorem 3.7] we know that $A' = A \otimes_R R'$ is solid for every Noetherian complete local domain R' and 1.6 gives that A' is parasolid. $\square$

Example 1.8. Let $R = K[[X, Y, Z]]/(XY, XZ)$, so that its spectrum is the union of an affine plane and an affine line meeting in one point. Let $A = K[[X, Y, Z]]/(X)$. Then A is a parasolid R -algebra, but not universally parasolid.

The following proposition characterizes parasolid R -algebras A for low dimensions (≤ 2) by $H_m^d(A) \neq 0$. Note that the characterization is set-theoretical for $d = 0$, topological for $d = 1$ and algebro-geometrical for $d = 2$.

Proposition 1.9. *Let R denote a local Noetherian ring of dimension d with maximal ideal $\mathfrak{m}$ and let A denote an R -algebra. Then the following are equivalent.*

$d = 0$.

- (i) A is parasolid.
- (ii) $A \neq 0$.
- (iii) $\text{Spec } A \neq \emptyset$.

$d = 1$. Let $x \in R$ be such that $V(x) = V(\mathfrak{m})$.

- (i) A is parasolid.
- (ii) $H_{\mathfrak{m}}^1(A) \neq 0$.
- (iii) $A \rightarrow A_x$ is not surjective (this means that $x^k \notin (x^{k+1})$ in A for all $k \in \mathbb{N}$.)
- (iv) $D(x) = D(\mathfrak{m}A) \subseteq \text{Spec } A$ is not a closed subset.

$d = 2$.

- (i) A is parasolid.
- (ii) $H_{\mathfrak{m}}^2(A) \neq 0$.
- (iii) $D(\mathfrak{m}A) \subseteq \text{Spec } A$ is not affine.

Proof. Suppose $d = 0$. For every R -module M we have

$$H_{\mathfrak{m}}^0(M) = \Gamma_{\mathfrak{m}}(M) = \{v \in M : \exists n \in \mathbb{N} \text{ such that } \mathfrak{m}^n v = 0\} = M.$$

In particular $H_{\mathfrak{m}}^0(R) = R$ and $1 \in R$ is the only paraclass, and it vanishes in $H_{\mathfrak{m}}^0(A)$ if and only if $A = 0$. (ii) $\Leftrightarrow$ (iii) is clear.

Suppose $d = 1$. (i) $\Rightarrow$ (ii) is clear. The cohomology sequence is

$$0 \longrightarrow H_{\mathfrak{m}}^0(A) \longrightarrow A \longrightarrow A_x \longrightarrow H_{\mathfrak{m}}^1(A) \longrightarrow 0$$

and this gives (ii) $\Leftrightarrow$ (iii). Since this is true for every parameter $x \in R$, we conclude (ii) $\Rightarrow$ (i). If $A \rightarrow A_x$ is surjective, then we have a closed embedding $D(x) \rightarrow \text{Spec } A$. On the other hand, consider $A \rightarrow A/\mathfrak{a} \rightarrow A_x$, where $\mathfrak{a}$ is the kernel. If $D(x)$ is closed, then $D(x) = V(\mathfrak{a})$, but x is a unit in $A/\mathfrak{a}$, hence $A/\mathfrak{a} \cong A_x$.

Suppose $d = 2$. (i) $\Rightarrow$ (ii) is again clear. If (ii) holds, then $H^1(D(\mathfrak{m}A), \mathcal{O}_A) \cong H_{\mathfrak{m}}^2(A) \neq 0$ and $D(\mathfrak{m}A)$ is not affine.

(iii) $\Rightarrow$ (i). Let x, y be parameters in R and assume that $c = 1/xy$ vanishes in $H_{\mathfrak{m}}^2(A) = H_{\mathfrak{m}A}^2(A)$. This means that there exists an equation

$$\frac{1}{xy} = \frac{a}{x^k} + \frac{b}{y^k}, \quad a, b \in A.$$

Hence we see that $q = b/y^{k-1} = 1/x - ay/x^k$ and $p = a/x^{k-1} = 1/y - bx/y^k$ are functions defined on $D(x, y) = D(\mathfrak{m}A) \subseteq \text{Spec } A$. Furthermore we see that

$$1 = \frac{a}{x^{k-1}}y + \frac{b}{y^{k-1}}x = px + qy,$$

and this shows that $D(x, y)$ is affine due to [4, Exercise II.2.17]. $\square$

2. ORDER IDEALS

Let A denote an R -algebra. An R -linear homomorphism $\varphi \in \text{Hom}_R(A, R)$ induce a mapping on the cohomology $H_{\mathfrak{m}}^d(\varphi) : H_{\mathfrak{m}}^d(A) \rightarrow H_{\mathfrak{m}}^d(R)$. These mappings are useful in the study of $H_{\mathfrak{m}}^d(A)$, in particular if R is complete.

Definition 2.1. Let A denote an R -algebra. The ideal

$$U(A) = \{\varphi(1) : \varphi : A \longrightarrow R \text{ } R\text{-linear}\},$$

is called the *order ideal* of the algebra A .

Remark 2.2. It is clear that this is an ideal and that an element $u \in R$ belongs to $U(A)$ if and only if u belongs to the image of some R -module homomorphism $A \rightarrow R$. The algebra A contains R as a direct summand if and only if $1 \in U(A)$. The algebra A over the domain R is solid if and only if $U(A) \neq 0$.

Let R be a local Noetherian ring with residue class field k and let E be an injective hull of k . Consider the Matlis-functor $D(-) = \text{Hom}_R(-, E)$. Let M denote an R -module. Then the mapping

$$H_{\mathfrak{m}}^d(M) \longrightarrow D(D(H_{\mathfrak{m}}^d(M)))$$

is injective due to [1, Remarks 10.2.2]. This means that for an element $0 \neq c \in H_{\mathfrak{m}}^d(M)$ there exists an R -morphism $\psi : H_{\mathfrak{m}}^d(M) \rightarrow E$ such that $\psi(c) \neq 0$.

If R is complete, then every R -module homomorphism $E \rightarrow E$ is the multiplication by a scalar $r \in R$, see [1, Theorem 10.2.12]. If furthermore R is Gorenstein, then $E = H_{\mathfrak{m}}^d(R)$ is an injective hull of the residue class field, see [1, Lemma 11.2.3].

Lemma 2.3. Let R denote a local complete Gorenstein ring of dimension d and let A be an R -algebra. Then the mapping

$$\text{Hom}_R(A, R) \longrightarrow \text{Hom}_R(H_{\mathfrak{m}}^d(A), H_{\mathfrak{m}}^d(R)), \quad \varphi \longmapsto H_{\mathfrak{m}}^d(\varphi)$$

is bijective.

Proof. We define an inverse mapping. Let $\psi : H_{\mathfrak{m}}^d(A) \rightarrow H_{\mathfrak{m}}^d(R)$ be R -linear and let $a \in A$. Consider

$$H_{\mathfrak{m}}^d(R) \xrightarrow{i} H_{\mathfrak{m}}^d(A) \xrightarrow{a} H_{\mathfrak{m}}^d(A) \xrightarrow{\psi} H_{\mathfrak{m}}^d(R).$$

The whole mapping is multiplication by an element $r \in R$, and we set $\varphi_{\psi}(a) := r$. This gives an R -linear mapping $\varphi_{\psi} : A \rightarrow R$.

Let $\varphi : A \rightarrow R$ be given. Then

$$(H_{\mathfrak{m}}^d(\varphi) \circ a \circ i)\left(\frac{s}{x_1 \cdots x_d}\right) = H_{\mathfrak{m}}^d(\varphi)\left(\frac{as}{x_1 \cdots x_d}\right) = \frac{\varphi(a)s}{x_1 \cdots x_d},$$

so this is multiplication by $\varphi(a)$.

Let $\psi : H_{\mathfrak{m}}^d(A) \rightarrow H_{\mathfrak{m}}^d(R)$ be given and let φ_{ψ} as just defined. Then

$$H_{\mathfrak{m}}^d(\varphi_{\psi})\left(\frac{a}{x_1 \cdots x_d}\right) = \frac{\varphi_{\psi}(a)}{x_1 \cdots x_d} = \psi\left(\frac{a}{x_1 \cdots x_d}\right).$$

□

Lemma 2.4. Let R denote a complete local Gorenstein ring of dimension d and let A be an R -algebra. Let $c \in H_{\mathfrak{m}}^d(R)$. Then the following are equivalent.

- (i) $i(c) \neq 0$ in $H_{\mathfrak{m}}^d(A)$.
- (ii) There exists a homomorphism $\varphi : A \rightarrow R$ such that $0 \neq H_{\mathfrak{m}}^d(\varphi)(i(c)) \in H_{\mathfrak{m}}^d(R)$.
- (iii) There exists an element $u \in U(A)$ such that $uc \neq 0$ in $H_{\mathfrak{m}}^d(R)$.
- (iv) $U(A) \not\subseteq \text{Ann}_R c$.

Proof. (i) $\Rightarrow$ (ii). From Matlis duality we know that there exists $\psi : H_{\mathfrak{m}}^d(A) \rightarrow H_{\mathfrak{m}}^d(R)$ such that $\psi(c) \neq 0$. Due to 2.3 this mapping is induced by $\varphi : A \rightarrow R$. Suppose (ii) holds. Then $u = \varphi(1)$ satisfies (iii). From (iii) to (i) is clear, and (iii) and (iv) are the same. $\square$

Corollary 2.5. *Let R denote a complete local Gorenstein ring of dimension d and let A be an R -algebra. Then the following are equivalent.*

- (i) A is parasolid.
- (ii) $U(A)$ is contained in no annihilator of a paraclass.

Proof. Follows from 2.4. $\square$

There exists always a class $c \in H_{\mathfrak{m}}^d(R)$ (in general no paraclass) such that its annihilator is $\text{Ann}_R c = \mathfrak{m}$.

Corollary 2.6. *Let R denote a complete local Gorenstein ring and let $c \in H_{\mathfrak{m}}^d(R)$ be a class such that $\text{Ann } c = \mathfrak{m}$. Let A denote an R -algebra. Then the following are equivalent.*

- (i) R is a direct summand in A
- (ii) The mapping $i : H_{\mathfrak{m}}^d(R) \rightarrow H_{\mathfrak{m}}^d(A)$ is injective.
- (iii) $i(c) \neq 0$ in $H_{\mathfrak{m}}^d(A)$.
- (iv) $U(A) = R$.

Proof. (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii) are clear. (iii) $\Rightarrow$ (iv). Let $i(c) \neq 0$. From 2.4 we know that $U(A) \not\subseteq \text{Ann } c = \mathfrak{m}$, hence $U(A)$ contains a unit and then $U(A) = R$. (iv) $\Rightarrow$ (i) is clear. $\square$

Lemma 2.7. *Let R denote a local complete Gorenstein ring of dimension d and let A denote an R -algebra. Let $i : H_{\mathfrak{m}}^d(R) \rightarrow H_{\mathfrak{m}}^d(A)$ denote the mapping on the local cohomology. Let $u \in R$. Then the following are equivalent.*

- (i) $u \in U(A)$.
- (ii) $u \in \bigcap_{c \in H_{\mathfrak{m}}^d(R)} (\text{Ann}_R c : \text{Ann}_R i(c))$.
- (iii) u annihilates $\text{kern } i$.

Proof. Suppose that (i) holds, say $u = \varphi(1)$. Let $r \in R$ and suppose that $ri(c) = 0$ in $H_{\mathfrak{m}}^d(A)$. Then $0 = \varphi(ri(c)) = urc =$ and ur annihilates c .

Suppose that (ii) holds, and let $i(c) = 0$. Then $\text{Ann } i(c) = R$ and $\text{Ann}(c) : R = \text{Ann}(c)$, hence $u \in \text{Ann}(c)$.

Suppose that (iii) holds. We have an injection $H_{\mathfrak{m}}^d(R)/\text{kern } i \hookrightarrow H_{\mathfrak{m}}^d(A)$. Due to the condition multiplication on $H_{\mathfrak{m}}^d(R)$ by u factors through $H_{\mathfrak{m}}^d(R)/\text{kern } i \rightarrow H_{\mathfrak{m}}^d(R)$. Since $H_{\mathfrak{m}}^d(R)$ is an injective module for a Gorenstein ring, there exists also an extension $H_{\mathfrak{m}}^d(A) \rightarrow H_{\mathfrak{m}}^d(R)$. This mapping comes from a mapping $\varphi : A \rightarrow R$ and it must hold $\varphi(1) = u$. $\square$

Lemma 2.8. *Let R be a Noetherian local Cohen-Macaulay ring of dimension d and let $x_1, \dots, x_d$ be parameters. Then the annihilator of the paraclass $1/x_1 \cdots x_d$ is just $(x_1, \dots, x_d)$.*

Proof. Every system of parameters is a regular sequence ([2, Theorem 2.1.2]). Consider the exact sequence $0 \rightarrow R \xrightarrow{x_d} R \rightarrow R/(x_d) \rightarrow 0$. The last part of the long exact sequence is

$$0 \longrightarrow H_{\mathfrak{m}}^{d-1}(R/(x_d)) \longrightarrow H_{\mathfrak{m}}^d(R) \xrightarrow{x_d} H_{\mathfrak{m}}^d(R) \longrightarrow 0.$$

Here $1/x_1 \cdots x_{d-1} \mapsto 1/x_1 \cdots x_d$, and the R -annihilator of these elements is the same. Hence we do induction, where the beginning is clear, since $H_m^0(R) = R$ for R zero-dimensional and $0 = \text{Ann } R$. $\square$

The property that R is a direct summand of A is in general much stronger than the property that A is parasolid. For complete regular rings however these properties coincide. From this result we will deduce in section 4 that every ideal in a regular ring is parasolidly closed.

Theorem 2.9. *Let R be a complete regular local ring and let A be an R -algebra. Then A is parasolid if and only if R is a direct summand of A .*

Proof. Let $\mathfrak{m} = (x_1, \dots, x_d)$, where d is the dimension of R . Then the annihilator of the corresponding paraclasse is exactly the maximal ideal $\mathfrak{m}$. If A is parasolid, then due to 2.5 we know that $U(A) \not\subseteq \mathfrak{m}$, hence $U(A) = R$. $\square$

3. PARASOLID CLOSURE

Let R denote a commutative ring and let $f_1, \dots, f_n$ and h be elements in R . Then we call the R -algebra

$$A = R[T_1, \dots, T_n]/(f_1 T_1 + \dots + f_n T_n + h)$$

the *forcing algebra* for the elements (or datas) $f_1, \dots, f_n; h$. The forcing algebra A forces $h \in (f_1, \dots, f_n)A$ and every R -algebra $R \rightarrow B$ such that $h \in (f_1, \dots, f_n)B$ factors through A .

Definition 3.1. Let R denote a commutative ring, let $f_1, \dots, f_n \in R$ and $h \in R$ be elements. Let A denote the forcing algebra for these elements.

If R is local and Noetherian, then we say that h *belongs parasolidly* to $f_1, \dots, f_n$ if the forcing algebra A is parasolid.

We say that h *belongs universally parasolidly* to $(f_1, \dots, f_n)$ if the forcing algebra A is universally parasolid.

Lemma 3.2. *Let R be a local Noetherian ring of dimension d and let $f_1, \dots, f_n$ and h be elements. Then the following hold.*

- (i) *If $h \in (f_1, \dots, f_n)$, then it belongs parasolidly to $f_1, \dots, f_n$.*
- (ii) *If $(f_1, \dots, f_n) \subseteq (g_1, \dots, g_m)$ and if h belongs parasolidly to $f_1, \dots, f_n$, then it belongs parasolidly to $g_1, \dots, g_m$.*
- (iii) *If h belongs parasolidly to $f_1, \dots, f_n$, then also rh belongs parasolidly to $f_1, \dots, f_n$.*
- (iv) *Let $a \in (f_1, \dots, f_n)$. Then h belongs parasolidly to $f_1, \dots, f_n$ if and only if $a + h$ belongs parasolidly to them.*

The same is true for R any Noetherian ring if we replace parasolidly by universally parasolidly.

Proof. (i). Let $h = r_1 f_1 + \dots + r_n f_n$, $r_i \in R$. This defines a section $A \rightarrow R$, thus R is a direct summand in A , hence A is parasolid. (ii) Let $f_i = r_{i1} g_1 + \dots + r_{im} g_m$ for $i = 1, \dots, n$. This defines a mapping

$$R[S_1, \dots, S_m]/(g_1 S_1 + \dots + g_m S_m + h) \longrightarrow R[T_1, \dots, T_n]/(f_1 T_1 + \dots + f_n T_n + h)$$

by sending $S_j \mapsto \sum_i r_{ij} T_i$. This is well defined since

$$\sum_{j=1}^m g_j S_j + h = \sum_{j=1}^m g_j \left(\sum_{i=1}^n r_{ij} T_i \right) + h = \sum_{i=1}^n \left(\sum_{j=1}^m r_{ij} g_j \right) T_i + h.$$

Since the algebra on the right is parasolid, so is the algebra on the left.

(iii). Here we use the mapping

$$R[S_1, \dots, S_n]/(f_1 S_1 + \dots + f_n S_n + rh) \longrightarrow R[T_1, \dots, T_n]/(f_1 T_1 + \dots + f_n T_n + h)$$

which sends $S_i \rightarrow rT_i$.

(iv). Let $a = a_1 f_1 + \dots + a_n f_n$. Then we use the mapping

$$R[S_1, \dots, S_n]/(f_1 S_1 + \dots + f_n S_n + h) \longrightarrow R[T_1, \dots, T_n]/(f_1 T_1 + \dots + f_n T_n + a + h)$$

where $S_i \mapsto T_i + a_i$.

The corresponding statement for universally parasolidly follows. $\square$

Definition 3.3. Let R denote a local complete Noetherian domain and let $h \in R$ and $f_1, \dots, f_n \in R$ be elements. We say that h belongs to the *parasolid closure* of $f_1, \dots, f_n$ if there exists a chain of elements $f_{n+1}, \dots, f_m = h$ such that f_i belongs universally parasolidly to $f_1, \dots, f_n, \dots, f_{i-1}$ for $i = n+1, \dots, m$.

Lemma 3.4. Let R denote a local complete Noetherian domain and let $f_1, \dots, f_n$ be elements. The elements which belong to the parasolid closure of $f_1, \dots, f_n$ form an ideal, which depends only on the ideal $(f_1, \dots, f_n)$.

Proof. Let h belong to the parasolid closure. This means that there exists a chain $f_1, \dots, f_n, f_{n+1}, \dots, f_m = h$, where f_i belongs universally parasolidly to $f_1, \dots, f_n, \dots, f_{i-1}$ for $i = n+1, \dots, m$. This is also true if we replace h by rh due to 3.2(iii). Let g be another element which belongs also to the parasolid closure, and let $f_1, \dots, f_n, g_1, \dots, g_k = g$ be a corresponding chain. Then also $f_1, \dots, f_n, f_{n+1}, \dots, f_m, g_1, \dots, g_k$ is such a chain due to 3.2(ii), and then due to 3.2(iv) also $f_1, \dots, f_n, f_{n+1}, \dots, f_m, g_1, \dots, g_k + f_m = g + h$ is such a chain. $\square$

Definition 3.5. Let R denote a Noetherian ring and let $I \subseteq R$ denote an ideal.

If R is a local complete domain, then we call the ideal just defined the *parasolid closure* I^* of I .

For general R , we define I^* by the condition that $h \in I^*$ if and only if this is true for all $\hat{R}_{\mathfrak{m}}/\mathfrak{q}$, where $\mathfrak{m}$ is a maximal ideal $\mathfrak{m}$ of R and $\mathfrak{q}$ is a minimal prime of the completion $\hat{R}_{\mathfrak{m}}$ (These rings are called the local complete domains of R).

Proposition 3.6. Let R denote a Noetherian ring and let $I \subseteq R$ be an ideal, $I = (f_1, \dots, f_n)$. Let $h \in R$. Then the following hold.

- (i) If h belongs universally parasolidly to I , then $h \in I^*$.
- (ii) $I \subseteq I^*$
- (iii) If $I \subseteq J$, then $I^* \subseteq J^*$.
- (iv) The parasolid closure is persistent: If $h \in I^*$, then for every ring homomorphism $R \rightarrow S$, S Noetherian, it holds that $h \in (IS)^*$.
- (v) Suppose that R is a local complete domain. Then $I = I^*$ if and only if no $h \notin I$ belongs universally parasolidly to $(f_1, \dots, f_n)$.
- (vi) $(I^*)^* = I^*$
- (vii) $h \in I^*$ if and only if $h \in (IR_{\mathfrak{m}})^*$ for every localization at a maximal ideal (or at a prime ideal).
- (viii) $h \in I^*$ if and only if $h \in (IR/\mathfrak{q})^*$ for every minimal prime ideal $\mathfrak{q}$ of R .
- (ix) Suppose that R is local. Then $h \in I^*$ if and only if $h \in (I\hat{R})^*$.

Proof. (i) and (ii) are clear. In (iii) we may assume that R is a local complete domain and then this follows from 3.2(ii). (iv). Let $S \rightarrow S'$ be a complete local

domain of S . This mapping factors then through a complete local domain R' of R , and in R' there exists a chain for h as in the definition 3.3 and this gives the chain in S' . (v) is clear, since every chain must start with an element which belongs universally parasolidly to the ideal. (vi). Let $h \in (I^*)^*$ and let R' denote a local complete domain of R . Then $(I^*)R' \subseteq (IR')^*$ due to persistence and then $h \in (I^*R')^* \subseteq ((IR')^*)^*$. Hence we may assume that R is a local complete domain, and the statement is clear from the definition by chains. (vii),(viii) and (ix) are clear by definition. $\square$

Remark 3.7. A natural question here is: can we through away some of the burden in the definition of parasolid closure which we needed to obtain the basic properties listed in 3.6? This refers in particular to the chains in 3.3, to localization and completion in 3.5 and to the use of all local Noetherian rings in the definition of universally parasolid. As for the last point, we will show in section 8 that for K -algebras of finite type it is enough to consider only local algebras which are essentially of finite type. (A variant with some advantages is to consider only those local rings where the extended ideal of I is primary to the maximal ideal.)

Lemma 3.8. *Let K be a field and let R be a Noetherian K -algebra, let $f_1, \dots, f_n \in R$ and $h \in R$. Let $\varphi : R \rightarrow S$ be a finite extension such that $h \in (f_1, \dots, f_n)S$. Then $h \in (f_1, \dots, f_n)^*$.*

Proof. Due to 1.4 (iii) the finite algebra S is universally parasolid. $\square$

Remark 3.9. The statement in the last lemma is open in the case of mixed characteristic and equivalent to the monomial conjecture. If we would define a parasolid algebra by the condition that not the paraclases $c \neq 0$ do not vanish, but only these paraclases which stay $\neq 0$ in every finite extension, then the contraction property would also hold in mixed characteristic, but then we could not prove the regular property anymore.

Example 3.10. The parasolid closure of 0 consists of the nilpotent elements. One inclusion is clear from the definition which refers to domains. If h is not nilpotent, then there exists a $R \rightarrow R' \neq 0$ where h becomes a unit and then the forcing algebra $R'/(h) = 0$ is not parasolid.

Example 3.11. Let C denote a local three-dimensional regular ring with maximal ideal $\mathfrak{m} = (x, y, z)$. Let

$$R = C/(x^i + y^j - z^k), \quad i, j, k \geq 1.$$

x and y are then parameters in R , we consider the forcing algebra for x, y and z , hence

$$A = R[T_1, T_2]/(xT_1 + yT_2 - z).$$

We have then the equation

$$x^i + y^j = (xT_1 + yT_2)^k \quad (*) .$$

Suppose that $k > i$ (or $k > j$). From the last equation we can read off the containment $x^i(1 - x^{k-i}T_1^k) \in (y)$. Since $k > i$ we have $x^k \in (x^{i+1})$, hence we get $x^i \in (y, x^{i+1})$ and this shows that $D(x, y) \subset \text{Spec } A$ is affine. Then the forcing algebra is not parasolid (not even solid) and $z \notin (x, y)^*$.

Suppose that $k = i = j$. From the equation $(*)$ we see $x^k(1 - T_1^k) \in (y)$, $y^k(1 - T_2^k) \in (x)$ and $\binom{k}{r} x^r y^{k-r} T_1^r T_2^{k-r} \in (x^{r+1}, y^{k-r+1})$ for $1 \leq r \leq k-1$.

Suppose that $k \geq 2$ and that at least one of the coefficients $\binom{k}{r}$, $1 \leq r \leq k-1$ is a unit in R . Then again $x^n y^m \in (x^{n+1}, y^{m+1})$ for some n, m and $z \notin (x, y)^*$.

4. REGULAR RINGS

Theorem 4.1. *Let R be a regular local ring and let $I \subseteq R$ be an ideal. Then $I = I^*$.*

Proof. We may assume that R is complete. Let $I = (f_1, \dots, f_n)$ and let $h \in R$ and let A denote the forcing algebra. Suppose that h belongs universally parasolidly to $f_1, \dots, f_n$. This means in particular that A is parasolid. Hence due to 2.9 R is a direct summand of A and then $h \in (f_1, \dots, f_n)$. $\square$

We give a second (more constructive) proof for the case that R contains a field. Again we may assume that R is complete, hence $R = K[[X_1, \dots, X_d]]$. Assume that $h \notin (f_1, \dots, f_n)$. Then also $h \notin (f_1, \dots, f_n, X_1^m, \dots, X_d^m)$ for $m \in \mathbb{Z}$ big enough and then we find $r = (r_1, \dots, r_d)$ such that $h \notin (I, X_1^{r_1+1}, \dots, X_d^{r_d+1})$, but $h \in (I, X_1^{r_1}, X_2^{r_2+1}, \dots, X_d^{r_d+1})$, $\dots$, $h \in (I, X_1^{r_1+1}, \dots, X_{d-1}^{r_{d-1}+1}, X_d^{r_d})$. In particular we have $hX_i \in (I, X_1^{r_1+1}, \dots, X_d^{r_d+1})$ for every variable X_i . We may replace I by $(I, X_1^{r_1+1}, \dots, X_d^{r_d+1}) = (f_1, \dots, f_n, X_1^{r_1+1}, \dots, X_d^{r_d+1})$ and we may assume that $f_j = \sum_{\nu \leq r} f_{j,\nu} X^\nu$ and also $h = \sum_{\nu \leq r} h_\nu X^\nu$.

There exists a K -linear form $\varphi : K[[X_1, \dots, X_d]] \rightarrow K$ given by $X^\sigma \mapsto c_\sigma$ such that $\varphi(I) = 0$, but $\varphi(h) = 1$. Modulo $(X_i^{r_i+1})$ we have $X^\sigma f_j = \sum_{\mu \leq r} f_{j,\mu-\sigma} X^\mu \in I$, hence $\sum_{\mu \leq r} c_\mu f_{j,\mu-\sigma} = 0$ (where $f_{j,\lambda} = 0$ if some entry of λ is negative). Furthermore $X^\sigma h = \sum_{\mu \leq r} h_{\mu-\sigma} X^\mu$ belongs also to I for $\sigma > 0$, hence $\sum_{\mu \leq r} c_\mu h_{\mu-\sigma} = 0$ for $\sigma > 0$, while $\sum_{\mu} c_\mu h_\mu = 1$. We consider

$$\sum_{\sigma \leq r} c_{r-\sigma} X^\sigma (f_1 T_1 + \dots + f_n T_n + h) = \sum_{\nu} a_\nu X^\nu,$$

where $a_\nu \in K[[T_1, \dots, T_n]]$ is the coefficient for X^ν . Then for $\nu \leq r$ we may write

$$\begin{aligned} a_\nu &= \sum_{\sigma} c_{r-\sigma} (f_1 T_1 + \dots + f_n T_n + h)_{\nu-\sigma} \\ &= \sum_{\mu} c_\mu (f_1 T_1 + \dots + f_n T_n + h)_{\mu-(r-\nu)} \\ &= \left(\sum_{\mu} c_\mu f_{1,\mu-(r-\nu)} \right) T_1 + \dots + \left(\sum_{\mu} c_\mu f_{n,\mu-(r-\nu)} \right) T_n + \sum_{\mu} c_\mu h_{\mu-(r-\nu)}. \end{aligned}$$

Since $\sum_{\mu \leq r} c_\mu f_{j,\mu-(r-\nu)} = 0$ for $\nu \leq r$ and $\sum_{\mu} c_\mu h_{\mu-(r-\nu)} = 0$ for $\nu < r$ and $\sum_{\mu} c_\mu h_\mu = 1$ for $\nu = r$ we see that $a_\nu = 0$ for $\nu < r$ and $a_r = 1$. Hence we get

$$\sum_{\sigma} c_{r-\sigma} X^\sigma (f_1 T_1 + \dots + f_n T_n + h) = X_1^{r_1} \dots X_d^{r_d} + R$$

where $R \in (X_1^{r_1+1}, \dots, X_d^{r_d+1})$, which shows that the paraclass for $X_1, \dots, X_d$ vanishes in the forcing algebra. $\square$

As usual in a tight closure type theory, we make the following definition.

Definition 4.2. We call a Noetherian ring R *pararegular* if $I = I^*$ holds for every ideal $I \subseteq R$.

The Theorem tells us that regular local rings are pararegular. This is also true for regular rings in general, since a locally pararegular ring is pararegular (the convers is not clear). In connection with persistence we get the following two corollaries.

Corollary 4.3. *Let R be a Noetherian ring which is a pure subring (e. g. a direct summand) of a (para)regular ring S . Then R is pararegular.*

Proof. Let $I \subseteq R$ denote an ideal, and let $h \in I^*$. Then also $h \in (IS)^* = IS$, but $R \cap IS = I$. $\square$

Corollary 4.4. *Let R be a Noetherian ring and let $I \subseteq R$ be an ideal. Then the parasolid closure of I is contained in the regular closure and in the integral closure, $I^* \subseteq I^{\text{reg}} \subseteq \bar{I}$.*

Proof. Let $h \in I^*$ and let $R \rightarrow S$ be a regular local ring. Then $h \in (IS)^* = IS$, so by definition it lies inside the regular closure. $\square$

Of course one hopes that pararegular rings are reduced, normal and Cohen-Macaulay, but we can prove only partial results. The Cohen-Macaulayness of parasolid rings follows in equal characteristic from the comparison with tight closure, see section 9.

Lemma 4.5. *Let R denote a Noetherian domain and let $\tilde{R}$ denote its normalization. Let $0 \neq f \in R$. If $h \in (f)^*$, then $h/f \in \tilde{R}$. If R contains a field, then also the converse holds.*

Proof. Let $h \in (f)^*$. This is then true in all discrete valuation domains B of R and hence $h \in fB$, so $h/f \in B$. Then h/f lies in the intersection of all valuation domains and hence in the normalization.

If $h/f \in \tilde{R}$, then consider the inclusion $R \subseteq R[h/f] \subseteq \tilde{R}$ and we find $h \in (f)$ in the finite extension $R[h/f]$, which is universally parasolid if R contains a field. $\square$

Proposition 4.6. *Let R denote a Noetherian ring and suppose that R is pararegular. Then the following holds.*

- (i) R is reduced.
- (ii) If R is a domain and contains a field, then it is normal.

Proof. This follows from 3.10 and 4.5. $\square$

Example 4.7. Let $R = K[U, V, W]/(UV - W^2) = K[X^2, Y^2, XY] \subset K[X, Y] = S$. Then the cohomology class $c = W/UV$ maps to $1/XY$. Therefore $c \neq 0$, and its annihilator is the maximal ideal (U, V, W) . The class c is no paraclass in $H_{\mathfrak{m}}^2(R)$, but in $H_{\mathfrak{m}}^2(S)$. Let A denote an R -algebra where $i(c) = 0$. This means that we have an equation $W(UV)^k = T_1 U^{k+1} + T_2 V^{k+1}$ and B factors through the forcing algebra A for this equation. This gives on S the forcing algebra with equation $XY(X^2 Y^2)^k = T_1 X^{2(k+1)} + T_2 Y^{2(k+1)}$. We know that $H_{\mathfrak{m}}^2(A \otimes_R S) = 0$ due to 4.1 and 1.9 and hence this is also true for $H_{\mathfrak{m}}^2(A)$, since $A \subset A \otimes_R S$ is a direct summand.

5. SOLID CLOSURE AND PARASOLID CLOSURE

We recall the definition of solid closure. Let R denote a Noetherian ring and let $(f_1, \dots, f_n) = I \subseteq R$ be an ideal, $h \in R$. Then $h \in I^{sc}$ if and only if the forcing algebra A on $\hat{R}_{\mathfrak{m}}/\mathfrak{q}$ is solid, where $\mathfrak{m}$ runs through the maximal ideals of R and $\mathfrak{q}$ is a minimal prime in the completion $\hat{R}_{\mathfrak{m}}$.

Proposition 5.1. *Let R be a Noetherian ring and let $I \subseteq R$ be an ideal. Then $I^* \subseteq I^{sc}$.*

Proof. We may assume that R is a local complete domain. It is enough to show that an element h which belongs universally parasolidly to I belongs to the solid closure of I . The forcing algebra A for I, h is then universally parasolid and so in particular solid. $\square$

Proposition 5.2. *Let K denote a field of positive characteristic p and let R denote a Noetherian K -Algebra. Let $I = (f_1, \dots, f_n) \subseteq R$ denote an ideal. Then $I^* = I^{sc}$.*

Proof. We have already shown the direct inclusion, thus suppose that $h \in I^{sc}$. Since solid closure is persistent [5, Theorem 5.6], we know for every local Noetherian complete domain $R \rightarrow R'$ that $H_{\mathfrak{m}}^d(A') \neq 0$, where d is the dimension of R' and A' is the forcing algebra over R' . By 1.6 A' is parasolid. $\square$

We obtain the connection to tight closure in positive characteristic.

Corollary 5.3. *Let K denote a field of positive characteristic p and let R denote a Noetherian K -Algebra. Suppose that R is an algebra essentially of finite type over an excellent local ring or that the Frobenius endomorphism in R is finite. Then $I^* = I^* = I^{sc}$ holds for every ideal $I \subseteq R$.*

Proof. The equation $I^* = I^{sc}$ is due to [5, Theorem 8.6]. $\square$

Remark 5.4. The conditions imposed in the corollary imply that R has a completely stable weak test element for tight closure. Even without this condition it is true that the tight closure lies inside the solid closure (= parasolid closure).

Example 5.5. We have a look at the example of Roberts. Let K be a field of characteristic zero and consider in the polynomial ring $R = K[X, Y, Z]$ the elements X^3, Y^3, Z^3 and $X^2Y^2Z^2$. The forcing algebra is then

$$A = K[X, Y, Z][T_1, T_2, T_3]/(X^3T_1 + Y^3T_2 + Z^3T_3 + X^2Y^2Z^2).$$

This algebra is not parasolid, since the paraclass $1/XYZ$ does vanish in $H_{\mathfrak{m}}^3(A)$, and $X^2Y^2Z^2$ does not belong to $(X^3, Y^3, Z^3)^* = (X^3, Y^3, Z^3)$.

On the other hand, the computation of Roberts in [12] shows that the class $1/X^2Y^2Z^2$ does not vanish in $H_{\mathfrak{m}}^3(A)$ and so A is a solid algebra (at least over $\hat{R}$) and $X^2Y^2Z^2$ belongs to $(X^3, Y^3, Z^3)^{sc}$. This example also shows that some paraclasses may survive, while others do not.

6. ORDER IDEAL AND TEST IDEAL

An R -module homomorphism $A = R[T_1, \dots, T_n]/(f_1T_1 + \dots + f_nT_n + h) \rightarrow R$ is given by $T^\nu \mapsto r_\nu \in R$ such that for every ν the condition

$$f_1r_{\nu+e_1} + \dots + f_nr_{\nu+e_n} = r_\nu$$

holds, where e_i is a standard base vector. It is in general not easy to construct such a morphism. The elements r_0 of such module homomorphisms build the order ideal. We can use such order elements to compute tight closure.

Lemma 6.1. *Let R be a commutative ring and let A be a forcing algebra for the elements $f_1, \dots, f_n, h$. Let $u \in U(A)$. Then $uh \in (f_1, \dots, f_n)$.*

If furthermore R contains a field of positive characteristic p , then $uh^q \in I^{[q]}$ for all $q = p^e$.

Proof. Let $u = \varphi(1)$. We apply φ to the forcing equation and get

$$uh = \varphi(1)h = \varphi(h) = f_1\varphi(T_1) + \dots + f_n\varphi(T_n).$$

Under the Frobenius homomorphism the forcing equation becomes $f_1^q T_1^q + \dots + f_n^q T_n^q = h^q$. We apply again φ and get $uh^q = f_1^q \varphi(T_1^q) + \dots + f_n^q \varphi(T_n^q)$. $\square$

Remark 6.2. If R is a domain over a field of positive characteristic, and if $U(A) \neq 0$, then an element $u \neq 0$ can be taken to show that $h \in I^*$. It is not clear whether an element u such that $uh^q \in I^{[q]}$ for all $q = p^e$ does belong to the order ideal (in the complete case). The proof of [5, Theorem 8.6] seems to give only that a power of u is an order element.

The following ideal is a candidate for a test ideal for parasolid closure.

Definition 6.3. Let R denote a local Noetherian complete ring of dimension d . We set

$$U(R) := \bigcap_{A \text{ universally parasolid forcing algebra}} U(A).$$

and

$$V = \langle c \in H_m^d(R) : \exists \text{ universally parasolid forcing algebra } i : R \rightarrow A, i(c) = 0 \rangle.$$

Lemma 6.4. *Let R denote a local complete Gorenstein ring. Then $U(R) = \text{Ann } V$.*

Proof. This follows from 2.7. Let $u \in U(R)$ and let $c \in V$ such that c vanishes in the universally parasolid forcing algebra A . By definition $u \in U(A)$ and then $uc = 0$, and this holds also in the submodule generated by such c . On the other hand, suppose that $u \in \text{Ann } V$ and let $i : R \rightarrow A$ be a universally parasolid forcing algebra. Then $\text{kern } i \subseteq V$ and u annihilates the kernel, hence $u \in U(A)$. (This correspondence is true for every family of algebras A_i .) $\square$

Proposition 6.5. *Let R denote a local complete Gorenstein ring which contains a field of positive characteristic. Let T be the tight closure test ideal of R . Then $T = U(R)$.*

Proof. We know that $T = \text{Ann}_R(0_{H_m^d(R)}^*)$, where $0_{H_m^d(R)}^*$ is the tight closure of the submodule $0 \subseteq H_m^d(R)$, see [10, Proposition 4.1]. Due to 6.4 it is enough to show that $V = 0_{H_m^d(R)}^*$. We know $0_{H_m^d(R)}^* = \{s/(x_1 \cdots x_d)^k : s \in (x_1^k, \dots, x_n^k)^*\}$. The inclusion $0_{H_m^d(R)}^* \subseteq V$ is clear, since $s/(x_1 \cdots x_d)^k$ is zero in the forcing algebra for $s; x_1^k, \dots, x_d^k$ and this forcing algebra is (formally solid, hence) universally parasolid.

So suppose that $c = s/(x_1 \cdots x_d)^k$ vanishes in a universally parasolid algebra B . This means that there exists $m \geq 1$ such that $s(x_1 \cdots x_d)^m \in (x_1^{k+m}, \dots, x_d^{k+m})$ in B . Then the forcing algebra A for the elements $s(x_1 \cdots x_d)^m; x_1^{k+m}, \dots, x_d^{k+m}$ maps to B , and since B is universally parasolid also A is universally parasolid.

Hence $s(x_1 \cdots x_d)^m \in (x_1^{k+m}, \dots, x_d^{k+m})^* = (x_1^{k+m}, \dots, x_d^{k+m})^*$ and this means that $s(x_1 \cdots x_d)^m / (x_1^{k+m} \cdots x_d^{k+m}) = s / (x_1^k \cdots x_d^k) \in 0^*$. Therefore we have $0_{H_m^d(R)}^* = V$. $\square$

7. PARAREGULAR RINGS

For a complete local Gorenstein ring there are a lot of different characterizations to be pararegular.

Proposition 7.1. *Let R denote a local complete Gorenstein ring of dimension d . Let $c \in H_m^d(R)$ be a class such that $\text{Ann } c = \mathfrak{m}$ (only in part (vii)). Then the following are equivalent.*

- (i) *R is pararegular, i.e. every ideal $I \subseteq R$ is parasolidly closed.*
- (ii) *Every universally parasolid forcing algebra A contains R as a direct summand.*
- (iii) *Every ideal generated by parameters is parasolidly closed.*
- (iv) *$V = 0$.*
- (v) *$U(R) = R$.*
- (vi) *Every universally parasolid R -algebra B contains R as a direct summand.*
- (vii) *$i(c) \neq 0$ in every universally parasolid R -algebra B .*

Proof. (i) and (ii) are equivalent for every local complete Noetherian ring due to 3.6 (v). (i) $\Rightarrow$ (iii) is clear.

(iii) and (iv) are equivalent for a local complete Cohen-Macaulay ring. Suppose that (iii) holds. Let $x_1, \dots, x_d$ be parameters and let $c = s/x_1 \cdots x_d$ and suppose that $i(c) = 0$ vanishes in the universally parasolid forcing algebra A . This means that $s(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ holds in A and then A factors through the forcing algebra to $x_1^{k+1}, \dots, x_d^{k+1}; s(x_1 \cdots x_d)^k$, which is then also universally parasolid. This means that $s(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})^* = (x_1^{k+1}, \dots, x_d^{k+1})$ in R and hence $c = 0$. For the converse suppose that s belongs universally parasolidly to $(x_1, \dots, x_d)$. Then $c = s/x_1 \cdots x_d$ is zero in the universally parasolid forcing Algebra, hence $c \in V = 0$. Then s annihilates the paraclass $1/x_1 \cdots x_d$ and since R is Cohen-Macaulay, it follows that $s \in (x_1, \dots, x_d)$.

Suppose furthermore that R is a complete Gorenstein ring. The equivalence (iv) $\Leftrightarrow$ (v) follows from 6.4.

(iii) $\Rightarrow$ (vi). Let $i : R \rightarrow B$ be a universally parasolid algebra. Due to 2.7 it is enough to show that the cohomological mapping $H_m^d(R) \rightarrow H_m^d(B)$ is injective. So let $c = s/x_1 \cdots x_d$ and suppose that $i(c) = 0$. This means that $s(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ holds in B and then again $s(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})^* = (x_1^{k+1}, \dots, x_d^{k+1})$ in R , thus $c = 0$.

(vi) $\Rightarrow$ (vii) and (vi) $\Rightarrow$ (ii) are clear. (vii) $\Rightarrow$ (v). If A is a universally parasolid forcing algebra, then $i(c) \neq 0$, hence $U(A) \not\subseteq \text{Ann } c = \mathfrak{m}$, thus $U(A) = R$. $\square$

Corollary 7.2. *Let R denote a local Gorenstein ring. Then R is pararegular if and only if $\hat{R}$ is pararegular.*

Proof. $\Leftarrow$ is always true and follows from 4.3. The converse holds because of 7.1 (iii). $\square$

Remark 7.3. Since direct summands of regular rings are pararegular, we get a lot of pararegular but non-regular rings. If we work over a base field K , then all

the quotient singularities are pararegular, for example $K[x, y, z]/(xy - z^n)$. This equation yields also in mixed characteristic pararegular rings, as the following computation shows.

Example 7.4. Let C denote a local three-dimensional regular ring with maximal ideal $\mathfrak{m} = (x, y, z)$ and consider

$$R = C/(xy - z^n), \quad n \geq 1.$$

The elements x and y are now parameters in R . We want to show that R is pararegular. The annihilator of $c = z^{n-1}/xy \in H_{\mathfrak{m}}^2(R)$ is the maximal ideal (x, y, z) . Due to 7.1 we have to show that if c vanishes in an R -algebra A , then A is not universally parasolid, and it is enough to look at a forcing algebra A given by an equation

$$z^{n-1}(xy)^k = T_1 x^{k+1} + T_2 y^{k+1}, \quad k \geq 0.$$

This equation yields

$$1 = \frac{T_1 x^{k+1}}{z^{n-1}(xy)^k} + \frac{T_2 y^{k+1}}{z^{n-1}(xy)^k} = \frac{T_1 x}{z^{n-1}y^k} + \frac{T_2 y}{z^{n-1}x^k} \quad (*) .$$

The equations

$$\frac{T_1 z}{y^{k+1}} = \frac{T_1 x z}{xy^{k+1}} = \frac{T_1 x z}{z^n y^k} = \frac{T_1 x}{z^{n-1}y^k} = 1 - \frac{T_2 y}{z^{n-1}x^k} = 1 - \frac{T_2 z}{x^{k+1}}$$

show that $T_1 x/z^{n-1}y^k$ (and also $T_2 y/z^{n-1}x^k$) is a function defined on $D(x, y) \subseteq \text{Spec } A$. We take the $2n$ power of $(*)$ and get

$$1 = \left(\frac{T_1 x}{z^{n-1}y^k} + \frac{T_2 y}{z^{n-1}x^k} \right)^{2n} = P\left(\frac{T_1 x}{z^{n-1}y^k}\right)^n + Q\left(\frac{T_2 y}{z^{n-1}x^k}\right)^n,$$

where P and Q are polynomials in $T_1 x/z^{n-1}y^k$ and $T_2 y/z^{n-1}x^k$ and hence defined on $D(x, y)$. We write

$$\left(\frac{T_1 x}{z^{n-1}y^k} \right)^n = \frac{T_1^n x^n}{z^{(n-1)n}y^{kn}} = \frac{T_1^n x^n}{(xy)^{n-1}y^{kn}} = \frac{T_1^n}{y^{n(k+1)-1}} x.$$

Hence it is enough to show that $T_1^n/y^{n(k+1)-1}$ is defined on $D(x, y)$, for then we know that (x, y) is the unit ideal in the ring of global sections $\Gamma(D(x, y), \mathcal{O}_A)$ (and $D(x, y)$ is affine and A is not parasolid). From $T_1 x^{k+1} = z^{n-1}(xy)^k - T_2 y^{k+1}$ we get

$$\begin{aligned} (T_1 x^{k+1})^n &= (z^{n-1}(xy)^k - T_2 y^{k+1})^n = y^{kn}(z^{n-1}x^k - T_2 y)^n \\ &= y^{kn} \sum_{i=0}^n \binom{n}{i} x^{ki} z^{(n-1)i} (-T_2 y)^{n-i} \\ &= y^{kn} \sum_{i=0}^n \binom{n}{i} x^{ki} z^{n(i-1)} z^{n-i} (-T_2 y)^{n-i} \\ &= y^{kn} \sum_{i=0}^n \binom{n}{i} x^{ki} (xy)^{i-1} (-T_2 z y)^{n-i} \\ &= \frac{y^{kn} y^{n-1}}{x} \sum_{i=0}^n \binom{n}{i} x^{ki+i} (-T_2 z)^{n-i} = \frac{y^{(k+1)n-1}}{x} (x^{k+1} - T_2 z)^n \end{aligned}$$

and hence $T_1^n/y^{n(k+1)-1} = (x^{k+1} - T_2 z)^n/x^{n(k+1)+1}$, so that this function is defined.

In the preceeding example it would have been enough to consider only $k = 0$, as the following lemma shows.

Lemma 7.5. *Let R denote a local two-dimensional Gorenstein ring. Let $c = s/xy \in H_m^2(R)$ be an element such that $\text{Ann } c = \mathfrak{m}$. Then R is pararegular if and only if $D(x, y) \subseteq \text{Spec } R[T_1, T_2]/(xT_1 + yT_2 + s)$ is affine.*

Proof. We may assume that R is complete. Due to 7.1 it is enough to show that if $c = 0$ in an R -algebra B , then B is not universally parasolid, and for that it is enough to show that $D(x, y) \subseteq \text{Spec } B$ is affine. If $c = 0$ holds in B then this algebra factors through a forcing algebra for elements $x^{i+1}, y^{j+1}; sx^i y^j$ for some $i, j \geq 0$. We have a ring homomorphism

$$R[S_1, S_2]/(sx^i y^j + x^{i+1} S_1 + y^{j+1} S_2) \longrightarrow R[T_1, T_2]/(s + xT_1 + yT_2)$$

given by $S_1 \mapsto y^j T_1, S_2 \mapsto x^i T_2$. This mapping is an isomorphism on $D(x, y)$ (check locally on $D(x)$ and $D(y)$), hence if $D(x, y)$ is affine on the right, it is affine on the left (In fact $D(x, y) \subseteq \text{Spec } R[T_1, T_2]/(xT_1 + yT_2 + s)$ is an affine-linear bundle of rank one over $D(x, y) \subseteq \text{Spec } R$ and it is a geometric representation (as a torseur) of the class $c = s/xy \in H^1(D(x, y), \mathcal{O}_R) \cong H_m^2(R)$). $\square$

8. EXPRESSING DATAS OVER $\mathbb{Z}$

Suppose that a certain R -algebra is not parasolid. Then we can express this fact by finitely many equations over $\mathbb{Z}$ and from this we get a lot of other rings R' where the corresponding algebra is also not parasolid. This gives in particular relations between the situation in positive characteristic, zero characteristic and mixed characteristic. The following lemma is the most general version of this observation.

Lemma 8.1. *Let D denote a commutative ring and let B be a finitely generated D -algebra. Suppose that there exists a local Noetherian D -algebra R of dimension d such that $B \otimes_D R$ is not parasolid. Then there exists a finitely generated D -algebra $S = D[X_1, \dots, X_d, Y_1, \dots, Y_m]/\mathfrak{a}$ such that*

- (i) $S \subseteq R$ and $X_1, \dots, X_d$ are parameters in R .
- (ii) The cohomology class $1/X_1 \cdots X_d \in H_{(X_1, \dots, X_d)}^d(S)$ is not zero.
- (iii) The cohomology class $1/X_1 \cdots X_d \in H_{(X_1, \dots, X_d)}^d(B \otimes_D S)$ is zero.

Moreover, for every local Noetherian S -algebra R' such that $X_1, \dots, X_d$ become parameters with non-vanishing paraclass, the algebra $B \otimes_D R'$ is not parasolid.

Proof. Let B be given by $B = D[T_1, \dots, T_n]/(P_1, \dots, P_m)$. Since $B \otimes_D R$ is not parasolid, there exists parameters $x_1, \dots, x_d$ in R (with non vanishing paraclass) such that their paraclass does vanish in $B \otimes_D R = R[T_1, \dots, T_n]/(P_1, \dots, P_m)$. This means that $(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ or that we have an equation

$$(x_1 \cdots x_d)^k = G_1 x_1^{k+1} + \dots + G_d x_d^{k+1} + H_1 P_1 + \dots + H_m P_m$$

where $G_i, H_j \in R[T_1, \dots, T_n]$. Let $y_j \in R$ be all the coefficients of G_i and H_j together. Now take the finitely generated D -subalgebra $S := D[x_1, \dots, x_d, y_j] \subseteq R$.

(i) is clear by definition and (ii) is clear, since the paraclass maps to the paraclass in R . (iii). We have the inclusion $S[T_1, \dots, T_n] \subseteq R[T_1, \dots, T_n]$ and so the last equation holds also in $S[T_1, \dots, T_n]$, and hence $(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ holds in $B \otimes_D S = S[T_1, \dots, T_n]/(P_1, \dots, P_m)$.

If $S \rightarrow R'$ is a mapping to a local Noetherian ring where the images of the X_i are parameters with non-vanishing paraclass, then this class vanishes in $B \otimes_D R'$, showing that this algebra is not parasolid. $\square$

Corollary 8.2. *Let $D = \mathbb{Z}[F_1, \dots, F_n, H]$ and let $B = D[T_1, \dots, T_n]/(F_1T_1 + \dots + F_nT_n + H)$ be the universal forcing algebra. Let R be a local Noetherian ring and let $f_1, \dots, f_n, h \in R$ be elements so that R is a D -algebra via $F_i \mapsto f_i$, $H \mapsto h$. Suppose that the forcing algebra $B \otimes_D R$ over R is not parasolid and that the paraclass to the parameters $x_1, \dots, x_d \in R$ is $\neq 0$, but vanishes in the forcing algebra. Then there exists a finitely generated $\mathbb{Z}$ -subalgebra $S \subseteq R$,*

$$S = \mathbb{Z}[F_1, \dots, F_n, H, X_1, \dots, X_d, Y_j]/\mathfrak{a}$$

such that the cohomology class $1/X_1 \cdots X_d \in H_{(X_1, \dots, X_d)}^d(S)$ is not zero, but it is zero in $H_{(X_1, \dots, X_d)}^d(B \otimes_D S)$.

Proof. This follows from 8.1. $\square$

Remark 8.3. The previous lemma leads to the following finitely generated $\mathbb{Z}$ -algebras (for $d, n, m, k \in \mathbb{N}$)

$$S = \mathbb{Z}[F_1, \dots, F_n, H, X_1, \dots, X_d, C_\nu, C_{1\nu}, \dots, C_{d\nu}, |\nu| \leq m]/\mathfrak{a},$$

where the relations (the generators of $\mathfrak{a} = \mathfrak{a}_k$) have to be chosen in such a way that they yield the equation

$$(X_1 \cdots X_d)^k + X_1^{k+1} \sum_{|\nu| \leq m} C_{1\nu} T^\nu + \dots + X_d^{k+1} \sum_{|\nu| \leq m} C_{d\nu} T^\nu + \sum_{|\nu| \leq m} C_\nu T^\nu \left(\sum_{i=1}^n T_i F_i + H \right)$$

in the forcing algebra for F_i, H over S . This gives the relations (looking at the coefficients of each monomial T^ν)

$$(X_1 \cdots X_d)^k + C_{10} X_1^{k+1} + \dots + C_{d0} X_d^{k+1} + C_0 H$$

and

$$C_{1\nu} X_1^{k+1} + \dots + C_{d\nu} X_d^{k+1} + C_\nu H + F_1 C_{\nu-e_1} + \dots + F_n C_{\nu-e_n},$$

where $|\nu| \leq m$, e_i is the i -th unit vector and $C_{\nu-e_j} = 0$ for $\nu_j = 0$.

Corollary 8.2 may be applied to show that if some rings containing a field are pararegular, then also the corresponding rings in mixed characteristic are pararegular. This technique may be thought of as a kind of reduction to the field case.

Proposition 8.4. *Let $G \in \mathbb{Z}[X, Y, Z]$ be a polynomial of type $G = XG_1 + YG_2 + Z^n$, $n \geq 1$ and suppose that $R = \mathbb{Q}[X, Y, Z]/(G)$ is pararegular. Let C denote a three-dimensional regular ring with maximal ideal $\mathfrak{m} = (x, y, z)$. Then $R' = C/(G)$ is pararegular except for finitely many residue characteristics.*

Proof. R' is a local Gorenstein ring and the elements x and y are parameters. The annihilator of the class z^{n-1}/xy is the maximal ideal (x, y, z) . Due to 7.5 we have to show that a geometric realization of this class over $D(x, y)$ is affine. Let $B = \mathbb{Z}[X, Y, Z][T_1, T_2]/(G, XT_1 + YT_2 + Z^{n-1})$ be the forcing algebra over $\mathbb{Z}$.

Due to our assumption the open subset $D(X, Y) \subseteq \text{Spec } B_{\mathbb{Q}}$ is affine. This means that there exists an equation

$$(XY)^k + C_1 X^{k+1} + C_2 Y^{k+1} + C_3 (XT_1 + YT_2 + Z^{n-1}) = 0$$

in $B_{\mathbb{Q}}$. Taking into account the coefficients in C_i , this equation holds also in B_g , $0 \neq g \in \mathbb{Z}$, hence

$$D(X, Y) \subseteq \operatorname{Spec} \mathbb{Z}_g[X, Y, Z][T_1, T_2]/(G, XT_1 + YT_2 + Z^{n-1})$$

is affine. If the residue characteristic of R' does not divide g , then we have a factorization $\mathbb{Z}_g[X, Y, Z]/(G) \rightarrow R'$ and $D(x, y)$ is affine in the forcing algebra over R' . $\square$

Remark 8.5. The previous proposition may be applied to the equations $xy + z^n = 0$, $x^2y + y^k + z^2 = 0$, $x^2 + y^3 + z^4 = 0$, $x^3 + xy^3 + y^2 + z^2 = 0$, $x^2 + y^3 + z^5 = 0$. These equations define over $\mathbb{Q}$ a quotient singularity, hence they are pararegular. Hence we know by the proposition that these equations give also in mixed characteristic pararegular rings except for finitely many characteristics. If we want to have more information about the number $g \in \mathbb{Z}$ we need to compute an explicit representation of the unit, as in example 7.4.

This can be obtained in the following way. We have an explicit quotient map

$$\varphi : \mathbb{A}_{\mathbb{Q}}^2 \rightarrow \operatorname{Spec} R, \quad \varphi(u, v) = (\varphi_1(u, v), \varphi_2(u, v), \varphi_3(u, v)) = (x, y, z).$$

When we pull back the forcing algebra A for x, y, z^{n-1} to the regular ring $\mathbb{Q}[u, v]$ we get a forcing algebra B over a regular ring which has an easy structure, in particular we can find functions p', q' defined on $D(u, v) \subseteq \operatorname{Spec} B$ such that $p'u + q'v = 1$. The group acts also on $\operatorname{Spec} B$ and on the ring of global sections $\Gamma(D(u, v), \mathcal{O}_B)$ and so we find also invariant functions p, q such that $px + qy = 1$. This gives the equation in A showing that $D(x, y)$ is affine.

This method works in principle also in higher dimensions, but then it is not clear that we only have to look at one forcing algebra.

Another feature of corollary 8.2 is that it allows us to apply Hochster's finiteness theorem (see [2, Theorem 8.4.1]) to parasolid closure and so to reduce statements in characteristic zero to positive characteristic. With this method we may prove the Theorem of Briançon-Skoda in characteristic zero for parasolid closure.

Proposition 8.6. *Let R denote a Noetherian ring containing a field K and let $I = (f_1, \dots, f_n)$ be an ideal. Then $\overline{I^{n+w}} \subseteq (I^{w+1})^*$.*

Proof. We may assume that R is local. In positive characteristic this is a standard result for tight closure (see [9, Theorem 5.7]), so by 5.3 and 5.4 it is also true for parasolid closure. Lets assume that the characteristic of K is zero and assume that we have a counter example in the local ring $R \supseteq \mathbb{Q}$, i.e. $h \in \overline{I^{n+w}}$ but $h \notin (I^{w+1})^*$. This means in particular that the forcing algebra for $f_1, \dots, f_n, h$ is not parasolid over $R \rightarrow R'$, but since the condition on the integral closure is persistent we may assume that $R = R'$. Let $x_1, \dots, x_d \in R$ be parameters in R such that their paraclass vanishes in the forcing algebra.

We have to express the relevant datas in a finitely generated $\mathbb{Z}$ -algebra. The condition $h \in \overline{I^{n+w}}$ means that we have an equation

$$h^s + a_1 h^{s-1} + \dots + a_s = 0, \quad \text{where } a_i \in (I^{n+w})^i = I^{i(n+w)}.$$

This means that we may write $a_i = \sum_{\nu, |\nu|=i(n+w)} r_{\nu} f_1^{\nu_1} \cdots f_n^{\nu_n}$. Furthermore the vanishing of the paraclass has to be expressed as in 8.3. Hence everything may be expressed by the $\mathbb{Z}$ -algebra

$$\mathbb{Z}[F_1, \dots, F_n, H, R_{\nu} (|\nu| = i(n+w), i = 1, \dots, s), X_1, \dots, X_d, C_{1\nu}, \dots, C_{d\nu}, C_{\nu}]/\mathfrak{a},$$

and since this algebra has a solution of height d (with respect to $X_1, \dots, X_d$) in a local Noetherian ring over a field of characteristic zero, it has also a solution of height d over a field of positive characteristic, see [2, Theorem 8.4.1]. This yields a contradiction. $\square$

Remark 8.7. We take a look at the situation of colon-capturing, i.e. suppose that $hy_t \in (y_1, \dots, y_{t-1})$, where $y_1, \dots, y_t$ are parameters. We would like to prove that $h \in (y_1, \dots, y_{t-1})^*$. This is true again via tight closure in positive characteristic, but applying the method of the last proposition shows only that the corresponding forcing algebra is parasolid, whereas universally parasolid is not clear due to the fact that the condition in colon-capturing is not persistent. Nevertheless we will show that colon-capturing holds for parasolid closure in equal characteristic zero in the next section.

Our definition of parasolid closure refers to all local Noetherian rings in order to make it persistent. The following proposition describes an important situation where it is enough to consider only algebras essentially of finite type.

Proposition 8.8. *Let K be a field and let R denote a K -algebra of finite type. Let A be an R -algebra of finite type and suppose that $A' = A \otimes_R R'$ is parasolid for every $R \rightarrow R'$, where R' is local and essentially of finite type over K . Then A is universally parasolid, i.e. A' is parasolid for every Noetherian local ring R' .*

Proof. Suppose that $R \rightarrow R'$ is a Noetherian local ring of dimension d such that $A \otimes_R R'$ is not parasolid. Due to 8.1 there is a finitely generated R -subalgebra $S \subseteq R'$ containing elements $x_1, \dots, x_d \in S$ which are parameters in R' and which have the properties of the lemma.

S is also finitely generated over the field K and the superheight of $(x_1, \dots, x_d)$ is exactly d . We know due to a theorem of Koh [11] that the superheight equals the finite superheight, hence there exists a morphism essentially of finite type $S \rightarrow R''$ (R'' may be chosen to be the localization at a maximal ideal of S' , $S \rightarrow S'$ of finite type) such that $(x_1, \dots, x_d)$ are parameters in the local ring R'' . But then their paraclass does vanish in $H_{\mathfrak{m}''}^d(A'')$ and we have a contradiction. $\square$

Corollary 8.9. *Let K be a field and let R denote a K -algebra of finite type. Let $I \subset R$ be an ideal and let $h \in R$ be an element, and let A denote the forcing algebra for h, I . If A' is parasolid for every $R \rightarrow R'$ essentially of finite type over K , R' local, then $h \in I^*$.*

Proof. This follows directly from 8.8. $\square$

9. GENERIC PROPERTIES

We want to show that tight closure in equal characteristic zero is contained inside the parasolid closure. For this we have to consider a relative situation over a base scheme $\text{Spec } D$ and we have to compare the parasolid closure in the generic fiber with the parasolid closure in the special fibers. For a D -algebra S and a point $\mu \in \text{Spec } D$ we denote by $S_\mu = S \otimes_D \kappa(\mu)$ the ring of the fiber over μ . Later on D will be a finitely generated $\mathbb{Z}$ -domain (often $D = \mathbb{Z}$) so that the closed points have positive characteristic while the generic point has characteristic zero.

Lemma 9.1. *Let D be a Noetherian domain with generic point $\eta \in \text{Spec } D$. Let S be an domain of finite type over D such that the fibers are equidimensional of*

dimension d over a non-empty open subset. Let $R = S_\eta$ be the ring of the generic fiber, which is a d -dimensional domain of finite type over $Q(D)$. Let $\mathfrak{m}$ denote a maximal ideal of R and suppose that $x_1, \dots, x_d \in R$ are parameters in $R_\mathfrak{m}$. Let A be an R -algebra of finite type and suppose that the paraclass $1/x_1 \cdots x_d$ vanishes in $H_\mathfrak{m}^d(A)$.

Then there exists a non-empty open affine subset $D(g) \subseteq \text{Spec } D$, a prime ideal $\mathfrak{n} \subseteq S$ and an S -algebra B of finite type such that the following hold.

- (i) The elements $x_1, \dots, x_d$ are defined on S_g and they are primary to $\mathfrak{n} \subseteq S_g$.
- (ii) For every point $\mu \in D(g)$ the intersection $\text{Spec } S_\mu \cap V(\mathfrak{n})$ is non-empty and zero-dimensional, and for every closed point $P \in \text{Spec } S_\mu \cap V(\mathfrak{n})$ the $x_1, \dots, x_d$ are parameters in $(S_\mu)_P$.
- (iii) $A = B_\eta$.
- (iv) For every $\mu \in D(g)$ and every closed point $P \in \text{Spec } S_\mu \cap V(\mathfrak{n})$ the paraclass $1/x_1 \cdots x_d \in H_P^d((S_\mu)_P)$ vanishes in $H_P^d((B_\mu)_P)$.

Proof. During the proof the open subset $D(g)$ gets smaller and smaller. Of course there exists $0 \neq g \in D$ such that $x_1, \dots, x_d \in S_g$. There exists a unique prime ideal $\mathfrak{n} \subset S$ such that $\mathfrak{n}S_\eta = \mathfrak{m}$, $\mathfrak{n} \cap D = \{0\}$. We have $(x_1, \dots, x_d) \subseteq \mathfrak{n}$ in S_g and $\mathfrak{n}$ is a minimal prime over (x_i) . Let $\mathfrak{n}_i$ denote the other minimal primes over $(x_1, \dots, x_d)$. Since these do not survive in S_η , there exist elements $0 \neq g_i \in D$ and $g_i \in \mathfrak{n}_i$. Making g_i to units, we may assume that $(x_1, \dots, x_d) \subseteq \mathfrak{n}$ is the only minimal prime. This gives (i).

The mapping $V(\mathfrak{n}) \rightarrow \text{Spec } D$ is finite in the generic point, hence it is finite over an open non-empty subset, and $V(\mathfrak{n}) \cap \text{Spec } S_\mu$ is zero-dimensional and non-empty. If P is a closed point in this intersection, then the x_i generate up to radical the maximal ideal $\mathfrak{m}_P$ in $(S_\mu)_P$. Since the dimension of every component of the fiber $\text{Spec } S_\mu$ is d , the height of P in the fiber is d , hence the x_i are parameters.

The construction of B is clear.

Generically we have $(x_1 \cdots x_d)^k \in (x_1^{k+1}, \dots, x_d^{k+1})$ in $A = B_\eta$. The coefficients of this equation belong to B_g and hence this equation holds over an open neighbourhood. Then it holds for every B_μ over $D(g)$ and this shows that the paraclass $1/x_1 \cdots x_d \in H_P^d((B_\mu)_P)$ vanishes. $\square$

Remark 9.2. If D is a Hilbert ring so that the closed points in $\text{Spec } D$ are dense, then the condition that the fibers are equidimensional over an open non-empty subset is superfluous, since this follows from [3, Theorem 14.8] and the condition that S is a domain.

Corollary 9.3. Let D be a Noetherian domain with generic point $\eta \in \text{Spec } D$. Let S be a domain of finite type over D such that the fibers are equidimensional. Let B be an S -algebra. Suppose that the set $\{\mu \in \text{Spec } D : S_\mu \rightarrow B_\mu \text{ is parasolid}\}$ is dense in $\text{Spec } D$. Then $S_\eta \rightarrow B_\eta$ is parasolid in the generic point.

Proof. Suppose that B_η is not parasolid in the generic point. This means that there exists a maximal ideal $\mathfrak{m} \subseteq S_\eta = R$ and parameters $x_1, \dots, x_d \in R_\mathfrak{m}$ such that their paraclass vanish in $(B_\eta)_\mathfrak{m}$. We find $0 \neq s \in S$, $s \notin \mathfrak{m}$ such that $x_1, \dots, x_d \in S_s =: S'$. We may then work with S' instead of S . Due to 9.1 the x_i are parameters in the closed points of the fibers over an open non-empty subset and their paraclass vanishes in B_μ , hence B_μ is not parasolid over S_μ for an open non-empty subset, yielding a contradiction. $\square$

We obtain an even stronger result if the special points have positive characteristic.

Corollary 9.4. *Let D be a $\mathbb{Z}$ -domain of finite type and let S be a domain of finite type over D . Suppose that the fibers $\text{Spec } S_\mu$ are normal (or analytically irreducible). Let B be an S -algebra. Suppose that the set $\{\mu \in \text{Spec } D : S_\mu \rightarrow B_\mu \text{ is parasolid}\}$ is dense in $\text{Spec } D$. Then $S_\eta \rightarrow B_\eta$ is universally parasolid in the generic point $\eta \in \text{Spec } D$.*

Proof. First note that the residue class fields of the closed points in $\text{Spec } D$ have positive characteristic. If for a closed point μ the S_μ -algebra B_μ is parasolid, then it is also universally parasolid due to the assumptions on the fibers and 1.7 and 1.5. We have to show that $A = B_\eta$ is universally parasolid over $R = S_\eta$ and due to 8.8 it is enough to show that A' is parasolid over R' , where R' is of finite type over R . There exists also an S -algebra S' of finite type such that $R' = S'_\eta$. Then the S' -algebra $B' = B \otimes_S S'$ is again parasolid over a dense subset of $\text{Spec } D$ and then also the generic algebra is parasolid. $\square$

We apply these results to forcing algebras and to the parasolid closure of an ideal.

Proposition 9.5. *Let D denote a Noetherian domain with generic point η . Let S be a D -domain of finite type such that the fibers are equidimensional, and let $R = S_\eta$ be the ring of the generic fiber. Let $I \subseteq S$ be an ideal and $h \in S$ and let B be the forcing algebra for h and I . Then the following hold.*

- (i) *If the forcing algebra $A = B_\eta$ is not parasolid in the generic point, then there exist a non-empty open subset $D(g) \subseteq \text{Spec } D$ such that the forcing algebra B_μ is not parasolid over S_μ for $\mu \in D(g)$.*
- (ii) *Suppose that $h \notin I^*$. Then there exists a non-empty open subset $D(g) \subseteq \text{Spec } D$ such that the forcing algebra B_μ is not universally parasolid over S_μ , $\mu \in D(g)$ (If the characteristic of $\kappa(\mu)$ is positive, this means that $h \notin I^*$ in S_μ .)*

Proof. The first statement follows from 9.3, so consider (ii). Since $h \notin I^*$, the forcing algebra B_η is not universally parasolid, hence we know due to 8.9 that there exists a ring homomorphism $R \rightarrow R'$ of finite type to a domain R' and a maximal ideal $\mathfrak{m}' \subseteq R'$ such that A' is not parasolid over $R'_{\mathfrak{m}'}$. There exists an S -domain S' of finite type such that $R' = S'_\eta$. Part (i) shows that there exists a non-empty open subset $D(g) \subseteq \text{Spec } D$ such that B'_μ is not parasolid over S'_μ for $\mu \in D(g)$, hence B_μ is not universally parasolid over S_μ for $\mu \in D(g)$. $\square$

There are different ways to define tight closure in equal characteristic zero all referring to positive characteristic. We cite the definition from [9, Appendix, definition 3.1].

Definition 9.6. Let C be a locally excellent Noetherian ring containing $\mathbb{Q}$, let $h \in C$ and let $I = (f_1, \dots, f_n) \subseteq C$ be an ideal. Then h belongs to the tight closure of I , $h \in I^*$, if there exists a finitely generated $\mathbb{Z}$ -subalgebra $S \subseteq C$ with $h \in S$ and $f_1, \dots, f_n \in S$ such that $h_\kappa \in I_\kappa^*$ in S_μ for almost all $\kappa = \kappa(\mu)$, $\mu \in \text{Spec } \mathbb{Z}$.

Proposition 9.7. *Let $\mathbb{Z} \subseteq S$ be a $\mathbb{Z}$ -domain of finite type and let $I \subseteq S$ be an ideal, $h \in S$. Set $R = S_\mathbb{Q}$ and suppose that $h \notin I^*$ in R . Then there exists an open non-empty subset $U \subseteq \text{Spec } \mathbb{Z}$ such that $h \notin I^*$ for every closed point $\mu \in U$.*

Proof. This follows from 9.5 and 5.3. $\square$

Remark 9.8. This result means that if $h \in I_{\kappa(\mu)}^*$ holds for a dense (=infinite) subset of points μ with positive characteristic, then $h \in I_{\mathbb{Q}}^*$ in the generic point (of characteristic zero). This is in particular true if $h \in I_{\kappa(\mu)}^*$ holds for an open non-empty subset – and this is the definition of tight closure in characteristic zero. It is not clear whether $h \in I_{\kappa(\mu)}^*$ can be true for a dense set without being true for almost all $\kappa(\mu)$ (see [9, Appendix 1.8]).

Corollary 9.9. *Let K denote a field of characteristic zero and let C be a Noetherian locally excellent K -algebra and let $I \subseteq C$ be an ideal. Then $I^* \subseteq I^*$.*

Proof. Suppose that $h \in I^*$. We may express the relevant datas in a finitely generated $\mathbb{Z}$ -algebra $S \subset C$ such that $S_{\mathbb{Q}} \subseteq C$ and where $h_{\kappa} \in I_{\kappa}^*$ holds for almost all κ . We may assume that C and S are domains. Then due to 9.7 we have $h \in I^*$ in $S_{\mathbb{Q}}$ and then due to the persistence of parasolid closure this holds also in C . $\square$

Remark 9.10. It follows from 9.9 that also colon capturing (and again the Theorem of Briançon-Skoda) holds in equal characteristic zero for parasolid closure. Of course one would like to prove these facts also without referring to positive characteristic, and also for mixed characteristic.

REFERENCES

- [1] M. P. Brodman and R. Y. Sharp: “Local Cohomology,” Cambridge University Press, Cambridge, 1998.
- [2] W. Bruns and J. Herzog, “Cohen-Macaulay-Rings,” revised edition, Cambridge Univ. Press, Cambridge, UK, 1998.
- [3] D. Eisenbud, “Commutative Algebra with a View Toward Algebraic Geometry,” Springer-Verlag, New York, 1995.
- [4] R. Hartshorne, “Algebraic Geometry,” Springer-Verlag, New York, 1977.
- [5] M. Hochster, Solid closure, Contemp. Math. **159** (1994), 103-172.
- [6] M. Hochster, Tight closure, Big CM Algebras and Solid Closure, Contemp. Math. **159** (1994), 173-196.
- [7] M. Hochster and C. Huneke, Tight closure, in Commutative Algebra, Math. Sci. Research Inst. Publ. **15** (1989), Springer-Verlag, 305-324.
- [8] M. Hochster and C. Huneke, Tight closure, invariant theory, and the Briançon-Skoda theorem, J. Amer. Math. Soc **3** (1990), 31-116.
- [9] C. Huneke, “Tight Closure and Its Application,” AMS, 1996.
- [10] C. Huneke, Tight Closure, Parameter Ideals, and Geometry, in Six Lectures on Commutative Algebra, Birkhäuser, Basel, 1998.
- [11] J. Koh, Super Height of an Ideal in a Noetherian Ring, J. Algebra **116** (1988), 1-6.
- [12] P. Roberts, A Computation of Local Cohomology, Contemp. Math. **159** (1994), 351-356.

MATHEMATISCHE FAKULTÄT, RUHR-UNIVERSITÄT, 44780 BOCHUM, GERMANY
E-mail address: `brenner@cobra.ruhr-uni-bochum.de`